

Testimony of Dr. Johannes Lehmann
Cornell University, Ithaca, NY, USA

Before the
House Select Committee on
Energy Independence and Global Warming

Thursday, June 18, 2009

2175 Rayburn House Office Building
Washington, DC

Biochar for sustainable carbon sequestration and global soil enhancement

Chairman Markey, Members of the Committee:

Thank you for the opportunity to provide scientific information about biochar carbon sequestration for sustainable climate change mitigation and global soil enhancement. Biochar is a fine-grained charcoal-like material that is produced through the heating of biomass under air-deprived conditions. This process is called pyrolysis. A wide variety of organic matter sources can be used as a feedstock for this process, including residues from forests or crop production, from animal production (manures), and from green waste streams, such as yard wastes. Upon pyrolysis at relatively low temperatures of 300-600°C, the chemical properties of biomass carbon change to form structures that are much more resistant to microbial degradation in comparison to the original organic matter. Thus, materials that would rapidly release carbon dioxide and other potent greenhouse gases as they decompose, are transformed into a material that degrades much more slowly, thereby creating a long-term carbon sink (Figure 1). Such thermally altered material is about 1.5 to 2 orders of magnitude more stable in soils than uncharred organic matter.^{1,2} Biochar has mean residence times of several hundreds to several thousands of years in soils.^{3,4}




Figure 1: Schematic showing how biochar would achieve an increase of soil carbon stocks by decreasing the carbon dioxide return to the atmosphere.⁵

The mechanism behind carbon sequestration through biochar in soils is very straightforward because stabilization is to a large extent a function of its intrinsic chemical stability. This is in contrast to uncharred organic matter, where soil carbon accrual primarily relies on a range of interactions between the mineral matrix and the organic matter.^{6,7} Therefore, the level at which soil carbon stores saturate⁸ and cease to

sequester additional carbon is greater for biochar than for uncharred crop or forestry residues.

Biochar is a familiar substance in soil. Most soils already contain char that was generated during vegetation fires throughout the past several thousand years. These chars are estimated to make up several percent of total soil organic carbon worldwide,^{9,10} which, in turn, is about twice the size of the atmospheric carbon pool.¹¹ Biochar soil management increases the amount of such naturally existing chars, which have been found to provide beneficial health and productivity properties to soil.

Biochar production and its application to soil provide several additional important value streams beyond direct climate change mitigation. These include waste management, energy production, and soil improvement (Figure 2). As a waste management strategy, biochar can be produced from a variety of feedstocks that would otherwise constitute a financial and environmental liability.¹² For example, in agricultural regions with high phosphorus and nitrogen levels in the soils and water, animal manures could be pyrolysed as a waste management strategy to prevent eutrophication. In many situations, compost, landfill or animal manure operations often generate large amounts of methane and nitrous oxide. By pyrolysing materials such as lawn clippings or biomass from forest thinning for fire prevention, the production of these even more potent non-CO₂ greenhouse gases would be effectively mitigated at the same time as the carbon is sequestered in soil.

A second value stream arises from bioenergy generated during biochar production. Between 2 and 7 units of energy can be produced for each unit of energy invested during the life cycle of various biochar systems.¹³ Biochar production can be paired with local heat generation such as a system where poultry manure is pyrolysed on-farm to heat barns and the resulting biochar is applied to fields.

The third value stream is the improvement of soil quality upon biochar additions. Crop yields can be significantly increased in soils that have productivity constraints. These may arise from degradation of soil organic matter or years of nutrient extraction through cultivation. The resulting losses of agrochemicals such as fertilizer nutrients, herbicides and pesticides can be mitigated by biochar's ability to retain these compounds.^{14,15,16,17,18} Subsequently, fertilizer use efficiency is increased.¹⁹ In its ability to improve several key properties of soils, biochar is particularly effective not only because it delivers these values for a longer period of time, but also because it has a greater effect per unit of carbon added to soil.²⁰ Improved soil fertility also provides better resilience against climate change. Taken together, these three sources of value have the potential to enhance food and energy security while also combating climate change.




Figure 2: Biochar value streams including biomass use, energy generation and soil enhancement.²¹

Deliberate biochar additions to soils have a number of implications for carbon trading. Additionality can be demonstrated because biochar is currently not actively produced or added to soils to any appreciable extent (less than 1% global penetration). However, it would still be critical to determine what the carbon stocks and flows would have been under the baseline system and to ensure that no greenhouse gas “leakage” would occur. For example, the land-use effects of large-scale biofuel-style plantations²² for biochar production or the removal and use of crop residues necessary to protect soil from erosion would likely make such systems inappropriate for biochar production to achieve net carbon sequestration. Measurement and verification of biochar sequestration is facilitated by the fact that the amount of carbon added at any one time is easily measured or calculated, and does not need to accumulate over time. Verification of lasting sequestration is possible because biochars bear a chemical signature that can be distinguished from other organic matter in soils. Furthermore, sequestered biochar carbon would not be released to the atmosphere due to changes in land management, fires, or deforestation, making it a strong candidate as a reliable carbon sequestration agent, with a mean residence time of several hundred to thousands of years.

The national or global potential of biochar to help mitigate climate change is only theoretical at this point, because too few biochar systems exist at scale of implementation. Conservative modeling of the technical potential place biochar as an approach to contribute on the order of 1Gt carbon removals annually by 2050 (considering only limited biomass feedstock availability and only carbon sequestration impacts).²³ Such widespread adoption of biochar systems will require sustainability criteria, since the climate change mitigation value of biochar arises from several connected sources including energy and agriculture. The potential for climate mitigation is highly variable from one biochar system to the next due to different feedstocks, scales, and applications^{13,24} which requires careful evaluation. Biochar must be integrated into existing food production systems and not be an alternative to food production, make use

of already developed best-management practices such as no-tillage or conservation agriculture, and, for efficiency, build on residue collection systems that are already in place.

While few fully implemented modern biochar systems exist worldwide, the necessary engineering and science capacity is available to evaluate a diverse set of biochar systems at scale of implementation in the near term. In fact, biochar science has rapidly evolved even over the past 12 months.²⁵ Evaluation does not rely on a fundamental advance in science, but on the application and adaptation of existing science. The underlying technology is robust and sufficiently simple to make it applicable to many regions globally.

Current hurdles to implementation are: availability of pyrolysis units at sufficient maturity to allow all necessary research and development, and, as a direct consequence, a lack of demonstrated carbon trading activities; of sufficient development of best biochar practices at scale of implementation, including farm scale; and of demonstration of soil health benefits for the full spectrum of agroecosystems. The distributed nature of biochar systems and the potential for variability between systems create significant opportunities for sustainability, but also hurdles to widespread adoption, regulation, and financial viability.

Establishment of policies at national and international levels is required to remove hurdles to implementation and support full evaluation of biochar systems. Mechanisms for carbon trading that recognize soil carbon sequestration, including biochar sequestration, need to be put into place. Methodologies must include full life cycle accounting of emissions balances to deliver net climate benefits. The entire value chain of mitigation approaches must be recognized, to reward those activities that have multiple environmental and societal benefits. Biochar must not be an alternative to making dramatic reductions in greenhouse gas emissions immediately, but it may be an important tool in our arsenal for combating dangerous climate change.

Thank you, Mr Chairman.

Summary

What is biochar?

- Biochar is a fine-grained charcoal-like material produced through pyrolysis.
- Pyrolysis is the heating of biomass to temperatures of 300-600°C under air-deprived conditions.
- Through pyrolysis, the feedstock changes chemically to form structures that are much more resistant to microbial degradation than the original material.
- Many different sources of organic matter can be used as a feedstock for this process, including residues from forests or crop production, from animal production (manures), and from green waste streams, such as yard wastes.
- Biochar-like materials produced through forest fires are already a significant part of the global soil carbon cycle.

How does biochar sequester carbon?

- Because biochar is much more stable than other forms of biomass-derived carbon in soil, it remains in the soil for much longer.
- Biochar is 1.5-2 orders of magnitude more stable in soils than uncharred material and has mean residence times of hundreds to thousands of years.
- The “saturation point” for biochar additions to soil would be significantly greater compared to other additions from organic matter.

Why is biochar valuable?

- Biochar is a very stable form of carbon and can thus be used to sequester CO₂.
- Biochar can be made from waste materials, including those (e.g., manure or green wastes) that may otherwise produce even more potent non-CO₂ greenhouse gases.
- Biochar production results in energy generation, which can also be integrated into sustainable local-scale operations such as the heating of farm buildings.
- Biochar’s addition to soils can enhance soil fertility and retention of agrochemicals.

What do we need to know/do?

- The technology and scientific knowledge is ready to implement the necessary steps to thoroughly develop biochar systems at a meaningful scale.
- This will be necessary in order to understand biochar best practices, demonstrate field-scale soil health benefits for different agroecosystems.
- Soil carbon sequestration, including biochar carbon sequestration must be recognized under carbon trading schemes.
- Robust guidelines must be developed to ensure that any integration of biochar into carbon trading schemes is truly additional, sustainable, and does not result in the “leakage” of greenhouse gas emissions.
- Biochar must not be seen as a replacement for dramatic reductions in our greenhouse gas emissions.

References

1. Baldock, J. A. and Smernik, R. J. 2002. Chemical composition and bioavailability of thermally altered *Pinus resinosa* (Red pine) wood. *Organic Geochemistry* 33: 1093-1109.
2. Cheng C.H., Lehmann J., Thies J.E., and Burton S. 2008. Stability of black carbon in soils across a climatic gradient. *Journal of Geophysical Research* 113: G02027.
3. Kuzyakov Y., Friedel J.K., Stahr K., 2000. Review of mechanisms and quantification of priming effects. *Soil Biology and Biochemistry* 32: 1485-1498.
4. Lehmann J., Czimczik C., Laird D., and Sohi S. 2009. Stability of biochar in soil. In: Lehmann J., Stephen J. (eds.) *Biochar for Environmental Management: Science and Technology*. Earthscan Publ., London, pp. 183-205.
5. Lehmann J. 2007a. A handful of carbon. *Nature* 447: 143-144.
6. Oades J.M. 1988. The retention of organic matter in soils. *Biogeochemistry* 5: 35–70.
7. Sollins P., Homann P., and Caldwell B.A. 1996. Stabilization and destabilization of soil organic matter: mechanisms and controls. *Geoderma* 74: 65-105.
8. Six J., Conant R.T., Paul E.A., and Paustian K. 2002. Stabilization mechanisms of soil organic matter: Implications for C-saturation of soils. *Plant and Soil* 241: 155-176.
9. González-Pérez J.A., González-Vila F.J., Almendros G., and Knicker H. 2004. The effect of fire on soil organic matter – a review. *Environment International* 30: 855-870.
10. Krull E., Lehmann J., Skjemstad J., Baldock J., and Spouncer L. 2008. The global extent of black C in soils: is it everywhere? In: Hans G. Schröder (ed.) *Grasslands: Ecology, Management and Restoration*. Nova Science Publishers, Inc., pp 13-17.
11. Denman K.L., Brasseur G., Chidthaisong A., Ciais P., Cox P.M., Dickinson R.E., Hauglustaine D., Heinze C., Holland E., Jacob D., Lohmann U., Ramachandran S., da Silva Dias P.L., Wofsy S.C., and Zhang X., 2007. Couplings Between Changes in the Climate System and Biogeochemistry. In: *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change* [Solomon, S., D. Qin, M. Manning, Z. Chen, M. Marquis, K.B. Averyt, M. Tignor and H.L. Miller (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.
12. Lehmann J., Gaunt J., and Rondon M. 2006. Bio-char sequestration in terrestrial ecosystems - a review. *Mitigation and Adaptation Strategies for Global Change* 11: 403-427.
13. Gaunt J. and Lehmann J. 2008. Energy balance and emissions associated with biochar sequestration and pyrolysis bioenergy production. *Environmental Science and Technology* 42: 4152-4158.
14. Lehmann J., da Silva Jr. J.P., Steiner C., Nehls T., Zech W., and Glaser B. 2003. Nutrient availability and leaching in an archaeological Anthrosol and a Ferralsol of the Central Amazon basin: fertilizer, manure and charcoal amendments. *Plant and Soil* 249: 343-357.

15. Sheng G., Yang T., Huang M., and Yang K. 2005. Influence of pH on pesticide sorption by soil containing wheat residue-derived char. *Environmental Pollution* 134: 457-463.
16. Hiller E., Fargasova A., Zemanova L., and Bartal M. 2007. Influence of wheat ash on the MCPA immobilization in agricultural soils. *Bulletin of Environmental Contamination and Toxicology* 78: 345-348.
17. Novak J.M., Busscher W.J., Laird D.L., Ahmedna M., Watts D.W., and Niandou M.A.S. 2009. Impact of biochar amendment on fertility of a southeastern coastal plain soil. *Agronomy Journal* 174: 1-8.
18. Yu X.Y., Ying G.G. and Koonaka R. S. 2009. Reduced plant uptake of pesticides with biochar additions to soil. *Chemosphere*, published online, doi:10.1016/j.chemosphere.2009.04.001
19. Steiner C., Glaser B., Teixeira W.G., Lehmann J., Blum W.E.H. and Zech W. 2008. Nitrogen retention and plant uptake on a highly weathered central Amazonian Ferralsol amended with compost and charcoal. *Journal of Plant Nutrition and Soil Science* 171: 893-899.
20. Liang B., Lehmann J., Solomon D., Kinyangi J., Grossman J., O'Neill B., Skjemstad J.O., Thies J., Luizão F.J., Petersen J. and Neves E.G. 2006. Black carbon increases cation exchange capacity in soils. *Soil Science Society of America Journal* 70: 1719-1730.
21. Lehmann J. 2007b. Bio-energy in the black. *Frontiers in Ecology and the Environment* 5: 381-387.
22. Searchinger T., Heimlich R., Houghton R.A., Dong F., Elobeid A., Fabiosa J., Tokgoz S., Hayes D., and Yu T.-H. 2008. Use of U.S. croplands for biofuels increases greenhouse gases through emissions from land-use change. *Science* 319: 1238-
23. Amonette J., Lehmann J., and Joseph S. 2007. Terrestrial carbon sequestration with biochar: A preliminary assessment of its global potential. *EOS Transaction, s American Geophysical Union* 88(52), Fall Meet. Suppl., Abstract U42A-06.
24. Gaunt J. and Cowie A. 2009. Biochar, greenhouse gas accounting and emissions trading. In: Lehmann J., Stephen J. (eds.) *Biochar for Environmental Management: Science and Technology*. Earthscan Publ., London, pp. 318-340.
25. Lehmann J. and Stephen J. (eds.) 2009. *Biochar for Environmental Management: Science and Technology*. Earthscan Publ., London, 404p